

AD-A101 922

PURDUE UNIV LAFAYETTE IN DEPT OF STATISTICS

F/G 12/1

SELECTION PROCEDURES FOR A PROBLEM IN ANALYSIS OF VARIANCE. (U)

JUN 81 S S GUPTA, D HUANG

N00014-75-C-0455

NL

UNCLASSIFIED

NMS-81-22

AD-A101 922


END
DATE
FILED
8-81
DTIC

50
AD A101922

LEVEL

(12)

PURDUE UNIVERSITY



DTIC
ELECTE
JUL 24 1981
D
C

DEPARTMENT OF STATISTICS

DIVISION OF MATHEMATICAL SCIENCES

DTIC FILE COPY

DISTRIBUTION STATEMENT A
Approved for public release;
Distribution Unlimited

81 7 21 025

(12)

Selection Procedures for A
Problem in Analysis of Variance*

by

(10) Shanti S. Gupta Purdue University
and
Deng-Yuan Huang Academia Sinica, Taipei

9 Mimeograph Series #81-22

(14) MMS-81-22

12 49

Department of Statistics
Division of Mathematical Sciences
Mimeograph Series #81-22

DTIC
ELECTE
JUL 24 1981

(11) June 1981

(13) *This research was supported by the Office of Naval Research contract
N00014-75-C-0455 at Purdue University. Reproduction in whole or in part
is permitted for any purpose of the United States Government.

DISTRIBUTION STATEMENT A
Approved for public release;
Distribution Unlimited

291731 64

Selection Procedures For A
Problem in Analysis of Variance*

Shanti S. Gupta Purdue University
and

Deng-Yuan Huang Academia Sinica, Taipei

1. Introduction

For a completely randomized block design with one observation per cell, we express the observable random variables $X_{i\alpha}$ ($i = 1, \dots, k$; $\alpha = 1, \dots, n$) as

$$(1.1) \quad X_{i\alpha} = \mu + \beta_{\alpha} + \tau_i + \epsilon_{i\alpha}, \quad \sum_{i=1}^k \tau_i = 0,$$

where μ is the mean-effect, β_1, \dots, β_n are the block effects (nuisance parameters for the fixed effects model), τ_1, \dots, τ_k are the treatment effects, and $\epsilon_{i\alpha}$ are the error components. We assume that the errors within each block are jointly normally distributed.

We assume that the quality of a treatment is judged by the largeness of the τ_i 's. A 'population' π_i is called the best if τ_i is the largest. In general, it may be complicated to derive suitable tests for appropriate hypotheses, in which the experimenter may really be interested. We apply the subset selection approach (using certain basic hypotheses) and thus obtain more appropriate information regarding the treatments. A subset selection procedure is designed to select a subset so as to include the best population. Selection of any subset that contains the best is called a correct selection (CS). Roughly speaking, any two populations that are in the same selected subset, will be considered as "equivalently good". If all populations are selected, we claim that all treatments are homogeneous. In general, for achieving the objective of the experimenter, one should establish a suitable set of basic hypotheses. Depending on the objective one should proceed to consider different

*This research was supported by the Office of Naval Research contract N00014-75-C-0455 at Purdue University. Reproduction in whole or in part is permitted for any purpose of the United States Government.

ways of formulating the basic hypotheses. In this paper, we discuss a method based on subset selection rules for the purpose of making a claim of the type: $\tau_i = \tau^* > \tau_j + \Delta$ for all $i \in I$ and $j \in J$, where I and J form a partition of $\{1, 2, \dots, k\}$. The process of making such a claim will be called hypothesis identification. This is achieved by setting up certain basic hypotheses regarding the τ_i 's and using a subset selection procedure to test these basic hypotheses. It should be pointed out that in identifying an appropriate hypothesis, we assume that the constant Δ in the claim is specified by the experimenter, say, based on past experience. Associated with the tests of the basic hypotheses using a selection rule, there are error probabilities and the infimum of the probability of a correct selection for the rule employed. These are related to the power function of these tests. The sum of the average (over the basic hypotheses tested) of the error probabilities and one minus the infimum of the probability of a correct selection is called the identification risk. The main theorem of the paper discusses the derivation of an optimal selection rule in the sense of minimizing the identification risk. For a more general theory of multiple decisions from ranking and selection approach, one can refer to a recent monograph by Gupta and Huang (1981). A general survey of the entire field is provided in Gupta and Panchapakesan (1979).

Let \underline{Y} be a random observable vector with probability distribution depending upon a parameter $\underline{\tau}' = (\tau_1, \dots, \tau_k) \in \Omega$. Consider a family of hypotheses testing problems as follows:

$$(1.2) \quad H_0: \underline{\tau} \in \Omega_0 \quad \text{vs} \quad H_i: \underline{\tau} \in \Omega_i, \quad 1 \leq i \leq k,$$

where $\Omega_0 = \{\underline{\tau} | \tau_1 = \dots = \tau_k\}$ and $\Omega_i = \{\underline{\tau} | \tau_i > \max_{j \neq i} \tau_j\}$, $i = 1, 2, \dots, k$. A test

of the hypotheses (1.2) will be defined to be a vector $(\delta_1(\underline{y}), \dots, \delta_k(\underline{y}))$,

where the elements of the vector are ordinary test functions; when \underline{y} is observed

$$(1.3) \quad E_{\tau} \delta_j(\underline{Y}) \leq \gamma, \quad \tau \in \Omega_0, \quad 1 \leq i \leq k,$$

where γ is the upper bound on the error probabilities associated with the treatment effects.

For each i , ($1 \leq i \leq k$), we would like to have $\beta_i(\underline{\tau})$ large when $\underline{\tau} \in \Omega_i$ subject to (1.3). For $\underline{\tau} \in \Omega_i$, if we make $\beta_i(\underline{\tau})$ large, then $\beta_j(\underline{\tau})$ should be small for $j \neq i$.

It should be pointed out that in the formulation and proof of the optimal selection procedure, results from Neyman-Pearson theory are used.

2. Formulation of an Optimal Selection Procedure

Assume that

$$\underline{x}'_\alpha = (x_{1\alpha}, \dots, x_{k\alpha}),$$

$\alpha = 1, \dots, n$, are independently and identically distributed random vectors with the following distribution:

$$(2.1) \quad (2\pi\sigma^2)^{-1/2} k n |\Lambda|^{-1/2} \exp[-\frac{1}{2\sigma^2} (\underline{x} - \underline{\theta})' \Lambda^{-1} (\underline{x} - \underline{\theta})],$$

where $\underline{x}' = (x_{11}, \dots, x_{k1}; \dots; x_{1n}, \dots, x_{kn})$ and $\underline{\theta}' = (\theta_{11}, \dots, \theta_{k1}; \dots; \theta_{1n}, \dots, \theta_{kn})$.

$$\theta_{i\alpha} = \mu + \beta_{\alpha} + \tau_i, \quad i = 1, \dots, k; \quad \alpha = 1, 2, \dots, n \text{ and } \Lambda \text{ is a known positive definite}$$

kn x kn correlation matrix defined as follows:

Accession For	
NTIS GRA&I	
DTIC TAB	
Unannounced	
Justification	
By	
Distribution/	
Availability Codes	
Avall and/or	
Dist	Special

$$\Lambda = (\lambda_{ij})_{k \times k \times n}$$

$$= \begin{bmatrix} \Lambda_1 & 0 & \dots & 0 \\ 0 & \Lambda_1 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \Lambda_1 \end{bmatrix}, \quad \text{where}$$

$$\Lambda_1 = \begin{bmatrix} 1 & \lambda \\ \lambda & 1 \end{bmatrix}_{k \times k}$$

We rewrite the original model as the general linear model as follows:

$$\underline{X} = \underline{\theta} + \underline{\epsilon}, \quad \underline{\epsilon} \sim N(\underline{0}, \sigma^2 \Lambda).$$

Since we are interested in the difference between all pairs of τ_i 's, we transform the linear model to the following: For any i , let

$$\underline{Y}_i = C \underline{\tau}_i + \underline{\eta}, \quad \underline{\eta} \sim N(\underline{0}, \sigma^2 \Sigma_i),$$

where $\underline{\tau}_i = (\tau_{i1}, \dots, \tau_{ik})$, $\tau_{ij} = \tau_i - \tau_j$, $j \neq i$,

$$\underline{Y}_i' = (Y_{i11}, \dots, Y_{ik1}; \dots; Y_{i1n}, \dots, Y_{ikn})_{1 \times (k-1)n}$$

$$Y_{ij\ell} = X_{i\ell} - X_{j\ell}, \quad i \neq j; \quad i, j = 1, \dots, k; \quad \ell = 1, \dots, n,$$

$$\underline{Y}_i = A_i \underline{X}, \quad \underline{\eta} = A_i \underline{\epsilon}$$

$$A_i = \begin{bmatrix} A_{i1} & & & \\ & A_{i1} & & \\ & & \ddots & \\ 0 & & & A_{i1} \end{bmatrix}_{(k-1)n \times kn}$$

$$\Sigma_i = A_i \Lambda A_i' = \begin{bmatrix} A_{i1} \Lambda_1 A_{i1}' & & 0 \\ & \ddots & \\ 0 & & A_{i1} \Lambda_1 A_{i1}' \end{bmatrix} (k-1)nx(k-1)n,$$

$$A_{i1} = \begin{bmatrix} & & & i & & \\ -1 & 0 & \dots & 0 & 1 & 0 & \dots & 0 \\ 0 & -1 & 0 & \dots & 0 & 1 & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & \dots & \dots & 0 & -1 & 1 & 0 & \dots & 0 \\ 0 & \dots & \dots & 0 & 1 & -1 & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & \dots & \dots & 1 & 0 & \dots & 0 & -1 \end{bmatrix} \begin{matrix} \\ \\ \\ i-1 \\ i+1 \\ (k-1) \times k \end{matrix}$$

$$C' = [I, \dots, I]_{(k-1) \times (k-1)n}$$

where each of the identity matrix in C' is $(k-1) \times (k-1)$.

The maximum likelihood estimator of $\underline{\tau}_i$ is as follows:

$$\hat{\underline{\tau}}_i = (C' \Sigma_i^{-1} C)^{-1} C' \Sigma_i^{-1} \underline{y}_i.$$

Since,

$$A_{i1} \Lambda_1 A_{i1}' = (1-\lambda) \begin{bmatrix} 2 & & 1 \\ & \ddots & \\ 1 & & 2 \end{bmatrix}_{(k-1) \times (k-1)}$$

$$(A_{i1} \Lambda_1 A_{i1}')^{-1} = (1-\lambda)^{-1} \frac{1}{k} \begin{bmatrix} k-1 & & -1 \\ & \ddots & \\ -1 & & k-1 \end{bmatrix} = V_i$$

$$C' \Sigma_i^{-1} C = n (A_{i1} \Lambda_1 A_{i1}')^{-1} = \frac{n}{k(1-\lambda)} \begin{bmatrix} k-1 & & -1 \\ & \ddots & \\ -1 & & k-1 \end{bmatrix}$$

$$(C' \Sigma_i^{-1} C)^{-1} = \frac{1-\lambda}{n} \begin{bmatrix} 2 & & 1 \\ & \ddots & \\ 1 & & 2 \end{bmatrix}_{(k-1) \times (k-1)},$$

$$C' \Sigma_i^{-1} = [I \dots I] \begin{bmatrix} v_i & & 0 \\ & \ddots & \\ 0 & & v_i \end{bmatrix}$$

$$= [v_i, \dots, v_i]$$

$$(C' \Sigma_i^{-1} C)^{-1} C' \Sigma_i^{-1} = \frac{1-\lambda}{n} \begin{bmatrix} 2 & & 1 \\ & \ddots & \\ 1 & & 2 \end{bmatrix} [v_i, \dots, v_i]$$

$$= \frac{1}{n} [I, \dots, I].$$

Hence,

$$\begin{aligned} \hat{\tau}_i &= (C' \Sigma_i^{-1} C)^{-1} C' \Sigma_i^{-1} y_i \\ &= \frac{1}{n} \begin{bmatrix} \sum_{\ell=1}^n y_{i1\ell} \\ \vdots \\ \sum_{\ell=1}^n y_{ik\ell} \end{bmatrix} = \begin{bmatrix} y_{i1} \\ \vdots \\ y_{ik} \end{bmatrix} = \begin{bmatrix} \bar{x}_i - \bar{x}_1 \\ \vdots \\ \bar{x}_i - \bar{x}_k \end{bmatrix}, \end{aligned}$$

where $\bar{x}_i = \frac{1}{n} \sum_{j=1}^n x_{ij}$, $1 \leq i \leq k$.

The joint density of $y_{i11}, \dots, y_{ik1}; \dots; y_{i1n}, \dots, y_{ikn}$ is the following:

$$p_{\tau_i}(y_i) = (2\pi\sigma^2)^{-\frac{1}{2}k} |\Sigma_i|^{-\frac{1}{2}} \exp\left[-\frac{1}{2\sigma^2} (y_i - C\tau_i)' \Sigma_i^{-1} (y_i - C\tau_i)\right]$$

where

$$\Sigma_i = A_i \Lambda A_i' = (1-\lambda) \begin{bmatrix} J & & 0 \\ & \ddots & \\ 0 & & J \end{bmatrix}_{(k-1)n \times (k-1)n}$$

$$J = \begin{bmatrix} 2 & & 0 \\ & \ddots & \\ 0 & & 2 \end{bmatrix}_{(k-1) \times (k-1)}.$$

$$\Sigma_i^{-1} = \begin{bmatrix} v_i & & 0 \\ & \ddots & \\ 0 & & v_i \end{bmatrix}.$$

Now, we specify the Ω_i 's as follows (Note that this is a different specification from that given earlier):

$$\Omega_i = \{\tau_i | \tau_i \geq \max_{j \neq i} \tau_j + \Delta\sigma\}, \quad 1 \leq i \leq k,$$

and

$$\bar{\Omega} = \bigcup_{i=1}^k \Omega_i.$$

Assume that σ is known. Let

$$\Delta_i' = (\Delta\sigma, \dots, \Delta\sigma)_{1 \times (k-1)}, \quad i = 1, \dots, k, \quad \Delta > 0.$$

Thus

$$\begin{aligned} \frac{p_{\Delta_i}(y_i)}{p_0(y_i)} &= \exp \frac{1}{2\sigma^2} \{-(y_i - C_{\Delta_i})' \Sigma_i^{-1} (y_i - C_{\Delta_i}) + y_i' \Sigma_i^{-1} y_i\} \\ &= \exp \left\{ \frac{1}{2\sigma^2} \Delta_i' C' \Sigma_i^{-1} y_i - \frac{1}{2\sigma^2} \Delta_i' C' \Sigma_i^{-1} C \Delta_i \right\} \\ &= \exp \left\{ \frac{n\Delta}{(1-\lambda)k\sigma} (y_{i1} + \dots + y_{ik}) - \frac{1}{2\sigma^2} \Delta_i' C' \Sigma_i^{-1} C \Delta_i \right\}. \end{aligned}$$

Hence, we can rewrite

$$\begin{aligned} \frac{p_{\Delta_i}(y_i)}{p_0(y_i)} &\geq d' \quad \text{as} \\ y_{i1} + \dots + y_{ik} &\geq d''\sigma. \end{aligned}$$

Let a selection rule $\delta^0 = (\delta_1^0, \dots, \delta_k^0)$ be defined by

$$\delta_i^0(y_i) = \begin{cases} 1 & \text{if } p_{\Delta_i}(y_i) \geq d' p_0(y_i), \\ 0 & \text{otherwise} \end{cases},$$

such that

$$(2.2) \quad E_{\tau} \delta^0(y_i) = \tau, \quad \tau \in \Omega_0. \quad \text{Then}$$

δ^0 maximizes

$$(2.3) \quad \inf_{\bar{\Omega}} P(CS|\delta)$$

among all selection rules $\delta \in S(\gamma)$.

Note that $\delta_i^0(y_i)$ is also based on the maximum likelihood estimators $\hat{\tau}_i$ of τ_i . Since for any $\delta \in S(\gamma)$,

$$\tau \in \bar{\Omega} = \bigcup_{i=1}^k \Omega_i \text{ implies } \tau \in \Omega_i \text{ for some } i, \text{ thus}$$

$$\begin{aligned} P(CS|\delta) &= \int \delta_i(y_i) p_{\tau}(y_i) dv(y_i) \\ &\geq \min_{1 \leq i \leq k} \inf_{\tau \in \Omega_i} \int \delta_i(y_i) p_{\tau}(y_i) dv(y_i). \end{aligned}$$

We have

$$\inf_{\tau \in \bar{\Omega}} P(CS|\delta) = \min_{1 \leq i \leq k} \inf_{\tau \in \Omega_i} \int \delta_i(y_i) p_{\tau}(y_i) dv(y_i).$$

For any $\delta \in S(\gamma)$, it follows that

$$\int (\delta_i - \delta_i^0)(p_{\Delta_i} - dp_0) \leq 0$$

which implies

$$\int \delta_i^0 p_{\Delta_i} \geq \int \delta_i p_{\Delta_i}.$$

Since $\delta_i^0(y_i)$ is nondecreasing in y_i , hence

$$\begin{aligned} \inf_{\tau \in \bar{\Omega}} P(CS|\delta^0) &= \min_{1 \leq i \leq k} \int \delta_i^0(y_i) p_{\Delta_i}(y_i) dv(y_i) \\ &\geq \min_{1 \leq i \leq k} \int \delta_i(y_i) p_{\Delta_i}(y_i) dv(y_i) \\ &\geq \min_{1 \leq i \leq k} \inf_{\tau \in \Omega_i} \int \delta_i(y_i) p_{\tau}(y_i) dv(y_i) \\ &= \inf_{\theta \in \bar{\Omega}} P(CS|\delta). \end{aligned}$$

We rewrite δ^0 as follows:

$$\delta_i^0(y_i) = \begin{cases} 1 & \text{if } y_{i1} + \dots + y_{ik} \geq d''\sigma, \\ 0 & \text{otherwise} \end{cases}$$

Thus, the optimal subset selection rule is as follows:

$$\delta_i^0(\underline{x}) = \begin{cases} 1 & \text{if } \bar{x}_i \geq \frac{1}{k-1} \sum_{j \neq i} \bar{x}_j + d\sigma, \\ 0 & \text{otherwise} \end{cases},$$

where $d = \frac{d''}{k-1}$.

Now, we wish to determine d and n . We make the following transformation:

$$z_{ik} = [1 \dots 1]_{1 \times (k-1)} \begin{bmatrix} y_{i1} \\ \vdots \\ y_{ik} \end{bmatrix}, \text{ and}$$

$$\tau = \tau_{i1} + \dots + \tau_{ik} = (k-1)\tau_i - \sum_{j \neq i} \tau_j.$$

Since the distribution of

$$\hat{\underline{z}}_i = \begin{bmatrix} y_{i1} \\ \vdots \\ y_{ik} \end{bmatrix} = (C' \Sigma_i^{-1} C)^{-1} C' \Sigma_i^{-1} \underline{y}_i$$

is $(2\pi\sigma^2)^{-\frac{1}{2}k} |\Sigma_{1i}|^{-\frac{1}{2}} \exp[-\frac{1}{2\sigma^2} (\hat{\underline{z}}_i - \underline{z}_i)' \Sigma_{1i}^{-1} (\hat{\underline{z}}_i - \underline{z}_i)]$, where $\Sigma_{1i} = \frac{1-\lambda}{n} J$.

Then the distribution of Z_{ik} is

$$[2\pi\sigma^2(1-\lambda)k(k-1) \frac{1}{n}]^{-\frac{1}{2}} \exp[-\frac{n}{2\sigma^2(1-\lambda)k(k-1)} (z_{ik} - \tau)^2].$$

Hence

$$\begin{aligned} E_0 \delta_i^0(\underline{y}_i) &= P(Z_{ik} \geq d''\sigma) \\ (2.4) \quad &= \Phi\left[-\frac{d''\sqrt{n}}{\sqrt{(1-\lambda)k(k-1)}}\right] = \gamma, \end{aligned}$$

and

$$\begin{aligned}
& \inf_{I \in \bar{\Omega}} P_I(CS | \delta^0) \\
&= \min_{1 \leq i \leq k} \int \delta_i^0(y_i) p_{\Delta_i}(y_i) dv(y_i) \\
&= \min_{1 \leq i \leq k} P_{\Delta_i}(Z_{ik} \geq d''\sigma) \\
&= \min_{1 \leq i \leq k} P_{\Delta_i} \left(\frac{(Z_{ik} - (k-1)\Delta)\sqrt{n}}{\sqrt{(1-\lambda)k(k-1)}} \geq \frac{(d'' - (k-1)\Delta)\sqrt{n}}{\sqrt{(1-\lambda)k(k-1)}} \right)
\end{aligned}$$

$$(2.5) \quad = \Phi \left[- \frac{(d'' - (k-1)\Delta)\sqrt{n}}{\sqrt{(1-\lambda)k(k-1)}} \right] = P^*.$$

For given r , P^* , k , λ , and Δ , we can find d'' and the smallest number of blocks, n , to satisfy equations (2.4) and (2.5). Note that this n is also the minimum sample size for the case of one observation per cell in the completely randomized block design.

We rewrite (2.4) and (2.5) as

$$\Phi \left[- \frac{d\sqrt{n(k-1)}}{\sqrt{(1-\lambda)k}} \right] = \gamma$$

and

$$\Phi \left[- \frac{(d-\Delta)\sqrt{n(k-1)}}{\sqrt{(1-\lambda)k}} \right] = P^*.$$

Let z_{P^*} and z_γ represent the upper percentage points corresponding to P^* and γ , respectively of the standard normal distribution. Then we have

$$d = - \frac{z_\gamma \Delta}{z_{P^*} - z_\gamma},$$

and

$$n = \left\langle \frac{(1-\lambda)k(z_{P^*} - z_\gamma)^2}{(k-1)\Delta^2} \right\rangle,$$

where $\langle a \rangle$ is the smallest integer greater than or equal to a .

Summarizing the previous results, we obtain the following theorem.

Theorem: Under model (1.1) with the stated assumption on $\underline{\epsilon}_\alpha$, an optimal procedure for selecting a subset of the "best" or "worthwhile" treatments based on the observed data \underline{x} and satisfying the conditions (2.2) and (2.3) is: Select the population π_i with probability $\delta_i^0(\underline{x})$ given by

$$\delta_i^0(\underline{x}) = \begin{cases} 1 & \text{if } \bar{x}_i \geq \frac{1}{k-1} \sum_{j \neq i} \bar{x}_j + d\sigma, \\ 0 & \text{otherwise} \end{cases},$$

where the smallest values of d and n are given by

$$d = - \frac{z_Y \Delta}{z_{P^*} - z_Y},$$

and

$$n = \left\langle \frac{(1-\lambda)k(z_{P^*} - z_Y)^2}{(k-1)\Delta^2} \right\rangle.$$

Furthermore, we have established the following connection between the selection procedure and the hypothesis identification problem as follows:

If $\pi_{i_1}, \pi_{i_2}, \dots, \pi_{i_j}$ ($j < k$) are selected, we say that these populations are homogeneous and make the hypothesis identification

$$H_i^1: \tau_{i_1} = \dots = \tau_{i_j} \geq \max_{\substack{1 \leq l \leq k \\ l \notin \{i_1, \dots, i_j\}}} \tau_l + \Delta\sigma.$$

Note that the overall identification risk connected with this problem is $\leq \gamma + (1-P^*)$.

Remark: It should be pointed out that for some pairs (γ, P^*) , δ^0 may not select any population. This is to be interpreted as not identifying any one of the appropriate hypotheses.

We consider some special cases to provide an idea as to the appropriate identification of one of the hypotheses. For $\gamma = 0.05, \lambda = 0.5$ and $P^* = 0.95, 0.90, 0.80$; then

(i) $k = 2$,

$$H_0: \tau_1 = \tau_2, H_1^1: \tau_1 \geq \tau_2 + \Delta\sigma, H_2^1: \tau_2 \geq \tau_1 + \Delta\sigma.$$

In this case, for specified Δ -values, the smallest d and n needed for the optimal selection rule are given in the following table.

Δ	0.1	0.5	1.0	2.0
$d(0.95, 0.90, 0.80)$	0.05, 0.06, 0.07	0.25, 0.32, 0.33	0.50, 0.64, 0.66	1.00, 1.29, 1.33
$n(0.95, 0.90, 0.80)$	1089, 858, 620	44, 35, 25	11, 9, 7	3, 3, 2

(ii) $k = 3$,

$$H_0: \tau_1 = \tau_2 = \tau_3, H_1^1: \tau_1 \geq \max(\tau_2, \tau_3) + \Delta\sigma,$$

$$H_2^1: \tau_2 \geq \max(\tau_1, \tau_3) + \Delta\sigma, H_3^1: \tau_3 \geq \max(\tau_1, \tau_2) + \Delta\sigma,$$

$$H_4^1: \tau_1 = \tau_2 \geq \tau_3 + \Delta\sigma, H_5^1: \tau_1 = \tau_3 \geq \tau_2 + \Delta\sigma,$$

$$H_6^1: \tau_2 = \tau_3 \geq \tau_1 + \Delta\sigma.$$

For optimal selection rule, the minimum value of d and n are computed (for specified values of Δ) and given in the following table.

Δ	0.1	0.5	1.0	2.0
$d(0.95, 0.90, 0.80)$	0.05, 0.06, 0.07	0.25, 0.32, 0.33	0.50, 0.64, 0.66	1.00, 1.29, 1.33
$n(0.95, 0.90, 0.80)$	817, 644, 465	33, 26, 19	9, 7, 5	3, 2, 2

(iii) $k = 4$,

$$H_0: \tau_1 = \tau_2 = \tau_3 = \tau_4, H_1^1: \tau_1 \geq \max(\tau_2, \tau_3, \tau_4) + \Delta\sigma,$$

$$\begin{aligned}
H_2^1: \tau_2 &\geq \max(\tau_1, \tau_3, \tau_4) + \Delta\sigma, & H_3^1: \tau_3 &\geq \max(\tau_1, \tau_2, \tau_4) + \Delta\sigma, \\
H_4^1: \tau_4 &\geq \max(\tau_1, \tau_2, \tau_3) + \Delta\sigma, & H_5^1: \tau_1 = \tau_2 &\geq \max(\tau_3, \tau_4) + \Delta\sigma, \\
H_6^1: \tau_1 = \tau_3 &\geq \max(\tau_2, \tau_4) + \Delta\sigma, & H_7^1: \tau_1 = \tau_4 &\geq \max(\tau_2, \tau_3) + \Delta\sigma, \\
H_8^1: \tau_2 = \tau_3 &\geq \max(\tau_1, \tau_4) + \Delta\sigma, & H_9^1: \tau_2 = \tau_4 &\geq \max(\tau_1, \tau_3) + \Delta\sigma, \\
H_{10}^1: \tau_3 = \tau_4 &\geq \max(\tau_1, \tau_2) + \Delta\sigma, & H_{11}^1: \tau_1 = \tau_2 = \tau_3 &\geq \tau_4 + \Delta\sigma, \\
H_{12}^1: \tau_1 = \tau_2 = \tau_4 &\geq \tau_3 + \Delta\sigma, & H_{13}^1: \tau_1 = \tau_3 = \tau_4 &\geq \tau_2 + \Delta\sigma, \\
H_{14}^1: \tau_2 = \tau_3 = \tau_4 &\geq \tau_1 + \Delta\sigma.
\end{aligned}$$

For the optimal selection rule, the minimum value of d and n are computed (for specified values of Δ) and given in the following table.

Δ	0.1	0.5	1.0	2.0
$d(0.95, 0.90, 0.80)$	0.05, 0.06, 0.07	0.25, 0.32, 0.33	0.50, 0.64, 0.66	1.00, 1.29, 1.33
$n(0.95, 0.90, 0.80)$	726, 572, 413	30, 23, 17	8, 6, 5	2, 2, 2

Note that P^* is the probability of correct selection for the associated subset selection rule, while the error probability γ is controlled at 5 percent level. The identification risk is $0.05 + (1 - P^*)$. We can explain the cases described above as follows: for $k = 2$, if the selected subset contains π_i only, we identify H_i^1 , $i = 1, 2$; if it contains π_1 and π_2 , we identify H_0 . For $k = 3$, if the selected subset contains π_i only, we identify H_i^1 , $i = 1, 2, 3$; if it contains π_1 and π_2 , π_1 and π_3 , or π_2 and π_3 only, we identify H_4^1 , H_5^1 or H_6^1 , respectively. Similar discussion applies to the case $k = 4$.

Now, we discuss the case where σ^2 is unknown. For any i , the maximum likelihood estimators of $\underline{\tau}_i$ and σ^2 are:

$$\hat{\underline{\tau}}_i = (C' \Sigma_i^{-1} C)^{-1} C^{-1} \Sigma_i^{-1} \underline{y}_i = \begin{bmatrix} y_{i1} \\ \vdots \\ y_{ik} \end{bmatrix}$$

and

$$\hat{\sigma}^2 = \frac{1}{(k-1)(n-1)} \underline{y}_i' [\Sigma_i^{-1} - \Sigma_i^{-1} C (C' \Sigma_i^{-1} C)^{-1} C' \Sigma_i^{-1}] \underline{y}_i.$$

We know that $\hat{\sigma}^2$ and $\hat{\underline{\tau}}_i$ are independent and the distribution $f(s)$ of $s = \frac{\hat{\sigma}}{\sigma}$ is $\sqrt{\chi_p^2(s)}$ with $p = (k-1)(n-1)$.

As before, we define the selection rule as follows:

$$\varphi_i^0(\hat{\underline{\tau}}_i, \hat{\sigma}) = \begin{cases} 1 & \text{if } y_{i1} + \dots + y_{ik} \geq d_1 \hat{\sigma}, \\ 0 & \text{otherwise} \end{cases},$$

or

$$\varphi_i^0(\underline{x}, \hat{\sigma}) = \begin{cases} 1 & \text{if } \bar{x}_i \geq \frac{1}{k-1} \sum_{j \neq i} \bar{x}_j + \frac{d_1}{k-1} \hat{\sigma} \\ 0 & \text{otherwise} \end{cases}.$$

Conditionally, for an observed value of $\hat{\sigma}$, we can discuss the optimality as before. However, the constant d and n can be determined without any difficulty by (2.8) and (2.9). Since

$$E_{\underline{\tau}} \varphi_i^0(\hat{\underline{\tau}}_i, \hat{\sigma}) = \gamma, \quad \underline{\tau} \in \Omega_0$$

we get

$$(2.6) \quad \int \phi \left[-\frac{d_1 s \sqrt{n}}{\sqrt{(1-\lambda)k(k-1)}} \right] f(s) ds = \gamma,$$

and

$$(2.7) \quad \inf_{\Omega} P(CS|\varphi^0) \\ = \int \phi \left[-\frac{(d_1 s - (k-1)\Delta)\sqrt{n}}{\sqrt{(1-\lambda)k(k-1)}} \right] f(s) ds = p^*.$$

This gives

$$(2.8) \quad t \left[-\frac{d_1 \sqrt{n(n-1)}}{\sqrt{(1-\lambda)k}} ; (k-1)(n-1), 0 \right] = \gamma,$$

and

$$(2.9) \quad t \left[-\frac{d_1 \sqrt{n(n-1)}}{\sqrt{(1-\lambda)k}} ; (k-1)(n-1), \frac{\Delta \sqrt{n(k-1)}}{\sqrt{(1-\lambda)k}} \right] = p^*,$$

where $t(a; b, c)$ is the percentage point of the noncentral t with b degrees of freedom and the noncentrality parameter c .

Acknowledgment:

The authors wish to thank Professor S. Panchapakesan for a critical reading of this paper and for suggestions to improve the presentation of this paper.

References

- [1] Gupta, S. S. and Huang, D. Y. (1981). Multiple Statistical Decision Theory. Lecture Notes in Statistics (6), Springer-Verlag, New York.
- [2] Gupta, S. S. and Panchapakesan, S. (1979). Multiple Decision Procedures. John Wiley and Sons, New York.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER Mimeograph Series #81-22	2. GOVT ACCESSION NO. A707921	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Selection Procedures for a Problem in Analysis of Variance	5. TYPE OF REPORT & PERIOD COVERED Technical	
7. AUTHOR(s) Shanti S. Gupta and Deng-Yuan Huang	6. PERFORMING ORG. REPORT NUMBER Mimeo. Series #81-22	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Purdue University Department of Statistics West Lafayette, Indiana 47907	8. CONTRACT OR GRANT NUMBER(s) ONR N00014-75-C-0455	
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Washington, DC	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	12. REPORT DATE June 1981	
	13. NUMBER OF PAGES 16	
	15. SECURITY CLASS. (of this report) Unclassified	
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release, distribution unlimited.		
17. DISTRIBUTION STATEMENT (of abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES YES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Multiple decision rules, Completely randomized block design, Joint normal distribution, Hypothesis identification, Identification risk, Subset selection, Optimal selection procedures.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) → There are many situations in the analysis of variance where an experimenter would like to make comparisons among (and select the "best" set) the treatments. In this paper we study the problem where the data are based on a completely randomized block design. It is shown that the subset selection approach is a useful method to make appropriate "identification" among the hypotheses and the selected subset. We propose an optimal selection procedure which controls the error probabilities when all the parameters (treatments) are equal and which maximizes the infimum of the probability of a correct selection over some preference parameter.		

DD FORM 1473
1 JAN 73

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

space, simultaneously. Some examples are provided to illustrate the optimal subset selection rule and its interpretation in terms of the "identified" hypotheses.

UNCLASSIFIED